
AXIUMBILIC SINGULAR POINTS ON SURFACES IMMERSED IN \mathbb{R}^4 AND THEIR GENERIC BIFURCATIONS

R. GARCIA, J. SOTOMAYOR, AND F. SPINDOLA

ABSTRACT. Here are described the *axiumbilic* points that appear in generic one parameter families of surfaces immersed in \mathbb{R}^4 . At these points the ellipse of curvature of the immersion, Little [7], Garcia - Sotomayor [11], has equal axes.

A review is made on the basic preliminaries on axial curvature lines and the associated axiumbilic points which are the singularities of the fields of *principal*, *mean axial lines*, *axial crossings* and the quartic differential equation defining them.

The Lie-Cartan vector field suspension of the quartic differential equation, giving a line field tangent to the Lie-Cartan surface (in the projective bundle of the source immersed surface which quadruply covers a punctured neighborhood of the axiumbilic point) whose integral curves project regularly on the lines of axial curvature.

In an appropriate Monge chart the configurations of the generic axiumbilic points, denoted by E_3 , E_4 and E_5 in [11] [12], are obtained by studying the integral curves of the Lie-Cartan vector field.

Elementary bifurcation theory is applied to the study of the transition and elimination between the axiumbilic generic points. The two generic patterns E_{34}^1 and E_{45}^1 are analysed and their axial configurations are explained in terms of their qualitative changes (bifurcations) with one parameter in the space of immersions, focusing on their close analogy with the saddle-node bifurcation for vector fields in the plane [1], [10].

This work can be regarded as a partial extension to \mathbb{R}^4 of the umbilic bifurcations in Garcia - Gutierrez - Sotomayor [5], for surfaces in \mathbb{R}^3 . With less restrictive differentiability hypotheses and distinct methodology it has points of contact with the results of Gutierrez - Guiñez - Castañeda [3].

INTRODUCTION

In this work are described the axiumbilic singularities, at which the ellipse of curvature, as defined in Little [7] and Garcia - Sotomayor [11], has equal axes. The focus here are the axiumbilic points that appear generically in one parameter families of surfaces immersed in \mathbb{R}^4 . It can be regarded as an extension from \mathbb{R}^3 to \mathbb{R}^4 , as target spaces for immersed surfaces, and from umbilic to axiumbilic points as singularities, of results obtained by Gutierrez - Garcia - Sotomayor in [5]. It is also a continuation, in the direction of bifurcations of axiumbilic singularities, of the study of the structural stability of global axial configurations started in Garcia - Sotomayor [11].

An outline of the organization of this paper follows:

Section 1 deals with geometric preliminaries and a review of axial lines and axiumbilic points in order to define the *principal* and *mean curvature* configurations and their quartic differential equations.

The third author was partially supported by a doctoral fellowship of CAPES. The second author participated in the FAPESP Thematic Project 2008/02841-4 and has a fellowship CAPES PVNS at UNIFEI. The first and second authors are partially supported by Pronex/FAPEG/CNPq Proc. 2012 10 26 7000 803. The first and second authors are fellows of CNPq.

In Section 2, locally presenting a surface M immersed into \mathbb{R}^4 with a Monge chart, are studied the axiumbilic points and the transversality conditions in terms of which are defined the generic axiumbilic points are made explicit.

Section 3 establishes the axial principal and mean configurations in a neighborhood of generic axiumbilic points, denoted E_3 , E_4 and E_5 . This description uses the suspension of Lie-Cartan, giving rise to a line field tangent to a surface, which quadruply covers a punctured neighborhood of the axiumbilic point, and whose integral lines project regularly on the lines of axial curvature. This follows the approach of Garcia and Sotomayor in [11] and [12], chap. 8. After this review follow two subsections devoted to describe the behaviors of the axial lines near the axiumbilic points denoted E_{34}^1 and E_{45}^1 , which are the transversal transitions between the generic axiumbilic points.

In fact, the axiumbilic point E_{34}^1 (Figure 7) characterizes the transition between an axiumbilic point of type E_3 and one of type E_4 , which is explained by the variation of one parameter family in the space of immersions \mathcal{C}^r , $r \geq 5$ of a surface M into \mathbb{R}^4 (Proposition 11), in a first analogy with the saddle-node bifurcation of vector fields [1], [10].

The axiumbilic point E_{45}^1 (Figure 11) is characterized by the collision and subsequent elimination between one point of type E_4 and other of type E_5 . Here also, this bifurcation phenomenon is explained by means of a parameter variation in the space of immersions (Proposition 17), in a second analogy with the saddle-node bifurcations in the plane [1] [10].

Section 4 establishes the genericity of the axiumbilic bifurcations studied in this paper.

This work is related to the papers by Guñez-Gutiérrez [2] and Guñez-Gutiérrez-Castañeda [3] where a description, in class \mathcal{C}^∞ and in the context of quartic differential forms, of the points E_{34}^1 and E_{45}^1 (using the notation H_{34} and H_{45}), can be found.

Here was adopted a different approach, using the Lie-Cartan suspension as established in Garcia-Sotomayor [11], for immersions of class \mathcal{C}^r , $5 \leq r \leq \infty$. This leads to an interpretation of these points with less restrictive differentiability hypotheses and allows proofs with techniques closer to those of elementary bifurcation theory as in [1] and [10].

Section 5 closes the paper with related comments on its results and their connection with others found in the literature.

Acknowledgment. The authors are grateful to the referee for his/her careful reading and helpful style suggestions.

1. DIFFERENTIAL EQUATION OF AXIAL LINES

Let $\alpha : M \rightarrow \mathbb{R}^4$ be an immersion of class \mathcal{C}^r , $r \geq 5$, of an oriented smooth surface in \mathbb{R}^4 , with the canonical orientation. Assume that (x, y) is a positive chart of M and that $\{\alpha_x, \alpha_y, N_1, N_2\}$ is a smooth positive frame in \mathbb{R}^4 , where for $\mathfrak{p} \in M$, $\{\alpha_x = \partial\alpha/\partial x, \alpha_y = \partial\alpha/\partial y\}_{\mathfrak{p}}$ is the standard basis of $T_{\mathfrak{p}}M$ in the chart (x, y) and $\{N_1, N_2\}_{\mathfrak{p}}$ is an orthonormal basis of the normal plane $N_{\mathfrak{p}}M$.

In the chart (x, y) , the first fundamental form is expressed by

$$I_\alpha = \langle D\alpha, D\alpha \rangle = E dx^2 + 2F dx dy + G dy^2$$

where, $E = \langle \alpha_x, \alpha_x \rangle$, $F = \langle \alpha_x, \alpha_y \rangle$ and $G = \langle \alpha_y, \alpha_y \rangle$ and the second fundamental form is given by $II_\alpha = II_\alpha^1 N_1 + II_\alpha^2 N_2$ where II_α^i , $i = 1, 2$, is

$$II_\alpha^i := \langle D^2\alpha, N_i \rangle = e_i dx^2 + 2f_i dx dy + g_i dy^2,$$

with $e_i = \langle \alpha_{xx}, N_i \rangle$, $f_i = \langle \alpha_{xy}, N_i \rangle$ and $g_i = \langle \alpha_{yy}, N_i \rangle$.

The *mean curvature vector* is defined by $H = h_1N_1 + h_2N_2$ with

$$h_i = \frac{Eg_i - 2Ff_i + Ge_i}{2(EG - F^2)}.$$

For $v \in T_{\mathbf{p}}M$, the *normal curvature vector in the direction v* is defined by:

$$(1) \quad k_n = k_n(\mathbf{p}, v) = \frac{II_\alpha(v)}{I_\alpha(v)} = \frac{II_\alpha^1(v)}{I_\alpha(v)}N_1 + \frac{II_\alpha^2(v)}{I_\alpha(v)}N_2.$$

The image of k_n restricted to the unitary circle $S_{\mathbf{p}}^1$ of $T_{\mathbf{p}}M$ describes in $N_{\mathbf{p}}M$ an ellipse centered in $H(\mathbf{p})$, which is called *ellipse of curvature* of α at \mathbf{p} , and it will be denoted by $\varepsilon_\alpha(\mathbf{p})$. When $(e_1 - g_1)f_2 - (e_2 - g_2)f_1 \neq 0$, it is an actual non-degenerate ellipse, which can be a circle. Otherwise it can be a segment or a point. As $k_n|_{S_{\mathbf{p}}^1}$ is quadratic, the pre-image of each point of the ellipse is formed of two antipodal points on $S_{\mathbf{p}}^1$, and therefore each point of $\varepsilon_\alpha(\mathbf{p})$ is associated to a direction in $T_{\mathbf{p}}M$. Moreover, for each pair of points in $\varepsilon_\alpha(\mathbf{p})$ antipodally symmetric with respect to $H(\mathbf{p})$, it is associated two orthogonal directions in $T_{\mathbf{p}}M$, defining a pair of *lines* in $T_{\mathbf{p}}M$ [7], [8], [9].

Consider the function:

$$\begin{aligned} \|k_n - H\|^2 &:= \left[\frac{e_1dx^2 + 2f_1dxdy + g_1dy^2}{Edx^2 + 2Fdxdy + Gdy^2} - \frac{Eg_1 - 2Ff_1 + Ge_1}{2(EG - F^2)} \right]^2 \\ &+ \left[\frac{e_2dx^2 + 2f_2dxdy + g_2dy^2}{Edx^2 + 2Fdxdy + Gdy^2} - \frac{Eg_2 - 2Ff_2 + Ge_2}{2(EG - F^2)} \right]^2 \end{aligned}$$

For each $\mathbf{p} \in M$ in which $\varepsilon_\alpha(\mathbf{p})$ is not a circle, the points maximum and minimum of this function determine four points over the ellipse of curvature $\varepsilon_\alpha(\mathbf{p})$, which are their vertices, located at the large and small axes.




FIGURE 1. Ellipse of curvature $\varepsilon_\alpha(\mathbf{p})$ and lines of axial curvature

As illustrated in Figure 1, to the small axis AB is associated the crossing $A'A''B'B''$ and to the large axis CD is associated the crossing $C'C''D'D''$. Thus, for each $\mathbf{p} \in M$ at which the non-degenerate ellipse is not a circle or a point, *two crossings* are defined in $T_{\mathbf{p}}M$, one associated to the large axis and the other to the small axis of the ellipse of curvature. These *fields of 2-crossings* in M are called *fields of axial curvature*.

Outside the set \mathcal{U}_α of points at which the ellipse of curvature is a circle (i.e. has equal axes), called *axiumbilic points*, the lines and crossings are said to be *lines and crossings of axial*

curvature. Those related to the large (respectively small) axis of the ellipse of curvature are called *lines and crossings of principal (respectively mean) axial curvature*.

From the considerations above, the axial directions are defined by the equationm

$$Jac(\|k_n - H\|^2, I_\alpha) = 0$$

which has four solutions for $\mathbf{p} \notin \mathcal{U}_\alpha$ and is singular at $\mathbf{p} \in \mathcal{U}_\alpha$. According to [11] and [12], the *differential equation of axial lines* is given by:

$$(2) \quad a_4 dy^4 + a_3 dy^3 dx + a_2 dy^2 dx^2 + a_1 dy dx^3 + a_0 dx^4 = 0,$$

where

$$\begin{aligned} a_4 &= -4F(EG - 2F^2)(g_1^2 + g_2^2) + 4G(EG - 4F^2)(f_1 g_1 + f_2 g_2), \\ &+ 8FG^2(f_1^2 + f_2^2) + 4FG^2(e_1 g_1 + e_2 g_2) - 4G^3(e_1 f_1 + e_2 f_2) \end{aligned}$$

$$\begin{aligned} a_3 &= -4E(EG - 4F^2)(g_1^2 + g_2^2) - 32EFG(f_1 g_1 + f_2 g_2), \\ &+ 16EG^2(f_1^2 + f_2^2) - 4G^3(e_1^2 + e_2^2) + 8EG^2(e_1 g_1 + e_2 g_2) \end{aligned}$$

$$\begin{aligned} a_2 &= -12FG^2(e_1^2 + e_2^2) + 12E^2F(EG - 4F^2)(g_1^2 + g_2^2), \\ &+ 24EG^2(e_1 f_1 + e_2 f_2) - 24E^2G(f_1 g_1 + f_2 g_2) \end{aligned}$$

$$\begin{aligned} a_1 &= 4E^3(g_1^2 + g_2^2) + 4G(EG - 4F^2)(e_1^2 + e_2^2) \\ &+ 32EFG(e_1 f_1 + e_2 f_2) - 16E^2G(f_1^2 + f_2^2) - 8E^2G(e_1 g_1 + e_2 g_2), \end{aligned}$$

$$\begin{aligned} a_0 &= 4F(EG - 2F^2)(e_1^2 + e_2^2) - 4E(EG - 4F^2)(e_1 f_1 + e_2 f_2) \\ &+ -8E^2F(f_1^2 + f_2^2) - 4E^2F(e_1 g_1 + e_2 g_2) + 4E^3(f_1 g_1 + f_2 g_2). \end{aligned}$$

Proposition 1 ([11], [12]). Let $\alpha : M \rightarrow \mathbb{R}^4$ be an immersion of class C^r , $r \geq 5$, of an oriented and smooth surface. Denote the first fundamental form of α by

$$I_\alpha = Edx^2 + 2Fdx dy + Gdy^2$$

and the second fundamental form by:

$$II_\alpha = (e_1 dx^2 + 2f_1 dx dy + g_1 dy^2)N_1 + (e_2 dx^2 + 2f_2 dx dy + g_2 dy^2)N_2$$

where $\{N_1, N_2\}$ is an orthonormal frame.

i) The differential equation of axial lines is given by:

$$\begin{aligned} \mathcal{G} &= [a_0 G(EG - 4F^2) + a_1 F(2F^2 - EG)]dy^4 \\ &+ [-8a_0 EFG + a_1 E(4F^2 - EG)]dy^3 dx \\ &+ [-6a_0 GE^2 + 3a_1 FE^2]dy^2 dx^2 + a_1 E^3 dy dx^3 + a_0 E^3 dx^4 = 0, \end{aligned}$$

where

$$\begin{aligned} a_1 &= 4G(EG - 4F^2)(e_1^2 + e_2^2) + 32EFG(e_1 f_1 + e_2 f_2) \\ &+ 4E^3(g_1^2 + g_2^2) - 8E^2G(e_1 g_1 + e_2 g_2) - 16E^2G(f_1^2 + f_2^2) \end{aligned}$$

and

$$\begin{aligned} a_0 &= 4F(EG - 2F^2)(e_1^2 + e_2^2) - 4E(EG - 4F^2)(e_1 f_1 + e_2 f_2) \\ &+ 4E^3(f_1 g_1 + f_2 g_2) - 4E^2F(e_1 g_1 + e_2 g_2) - 8E^2F(f_1^2 + f_2^2). \end{aligned}$$

ii) The axiumbilic points of α are characterized by $a_0 = a_1 = 0$.

The axiumbilic points are defined by the intersection of the curves $a_0(x, y) = 0$ and $a_1(x, y) = 0$. Assume, with no loss of generality, that they intersect at $(x, y) = (0, 0)$. In this work it will be considered the case where the intersection is transversal or quadratic at $(0, 0)$.

Figure 2 illustrates the generic contact of the curves $a_0(x, y) = 0$ and $a_1(x, y) = 0$, whose intersection characterizes the axiumbilic points.




FIGURE 2. Transversal and quadratic contact between the curves $a_0 = 0$ and $a_1 = 0$ at an axiumbilic point \mathbf{p} .

An axiumbilic point given by $(x, y) = (0, 0)$ is called *transversal* if

$$(3) \quad \frac{\partial(a_0, a_1)}{\partial(x, y)} \Big|_{(0,0)} = \begin{vmatrix} \frac{\partial a_0}{\partial x}(0, 0) & \frac{\partial a_0}{\partial y}(0, 0) \\ \frac{\partial a_1}{\partial x}(0, 0) & \frac{\partial a_1}{\partial y}(0, 0) \end{vmatrix} \neq 0.$$

The axiumbilic point given by $(x, y) = (0, 0)$ is said to be of *quadratic type* if the matrix

$$(4) \quad \frac{\partial(a_0, a_1)}{\partial(x, y)} \Big|_{(0,0)} = \begin{bmatrix} \frac{\partial a_0}{\partial x}(0, 0) & \frac{\partial a_0}{\partial y}(0, 0) \\ \frac{\partial a_1}{\partial x}(0, 0) & \frac{\partial a_1}{\partial y}(0, 0) \end{bmatrix}$$

has rank 1 and, assuming $\frac{\partial a_0}{\partial y}(0, 0) \neq 0$, it follows from the implicit function theorem that $y(x)$ is a local solution of $a_0(x, y(x)) = 0$. Writing $s(x) = a_1(x, y(x))$ it follows that $s'(0) = 0$ and $s''(0) \neq 0$.

A similar analysis can be carried out if other element of the matrix $\frac{\partial(a_0, a_1)}{\partial(x, y)} \Big|_{(0,0)}$ is non zero.

Remark 2 ([11]). In isothermic coordinates, where $E = G$ and $F = 0$, it follows that

$$\begin{aligned} a_1 = -a_3 &= E^3[e_1^2 + e_2^2 + g_1^2 + g_2^2 - 4(f_1^2 + f_2^2) - 2(e_1g_1 + e_2g_2)] \\ a_0 = a_4 &= -\frac{a_2}{6} = 4E^3[f_1g_1 + f_2g_2 - (e_1f_1 + e_2f_2)] \end{aligned}$$

and the differential equation of axial lines is simplified to

$$(5) \quad a_0(x, y)(dx^4 - 6dx^2dy^2 + dy^4) + a_1(x, y)(dx^2 - dy^2)dxdy = 0.$$

1.1. **Axial configurations of immersed surfaces in \mathbb{R}^4 .** Let $\mathcal{I}^r = \mathcal{I}^r(M, \mathbb{R}^4)$ the set of immersions of class \mathcal{C}^r . For $\alpha \in \mathcal{I}^r$, the differential equation of axial lines is well defined (equation (2)):

$$(6) \quad \mathcal{G}(x, y, dx, dy) = a_4dy^4 + a_3dy^3dx + a_2dy^2dx^2 + a_1dydx^3 + a_0dx^4 = 0$$

in the projective bundle PM of M .

For each $\alpha \in \mathcal{I}^r$, define the *Lie-Cartan surface* of the immersion α by $\mathbb{L}_\alpha := \mathcal{G}_\alpha^{-1}(0)$, which is of class \mathcal{C}^{r-2} , regular in $M - \mathcal{U}_\alpha$ and may present singularities at \mathcal{U}_α .

Moreover, as the set defined by the quartic equation (6) contains the projective lines at \mathcal{U}_α , it follows that \mathbb{L}_α is a ramified covering of degree 4 in $M - \mathcal{U}_\alpha$ and contains the projective line $\pi^{-1}(\mathbf{p})$ for each $\mathbf{p} \in \mathcal{U}_\alpha$.

In the chart (x, y, p) of PM , with $p = \frac{dy}{dx}$, equation (6) is given by

$$(7) \quad \mathcal{G}(x, y, p) = a_4 p^4 + a_3 p^3 + a_2 p^2 + a_1 p + a_0 = 0.$$

Consider the *Lie-Cartan vector field* X_α , of class \mathcal{C}^{r-3} , tangent to the surface $\mathcal{G} = 0$

$$(8) \quad X_\alpha := \mathcal{G}_p \frac{\partial}{\partial x} + p \mathcal{G}_p \frac{\partial}{\partial y} - (\mathcal{G}_x + p \mathcal{G}_y) \frac{\partial}{\partial p}.$$

The *axial curvature lines* are the projections by $\pi : PM \rightarrow M$ restricted to \mathbb{L}_α , of the integral curves of X_α .

See illustration in Figure 3. For each $\mathbf{p} \in M - \mathcal{U}_\alpha$ there are 4 well defined axial directions, given the four roots of equation (7).

Two *axial configurations* are given: the *principal axial configuration* $\mathcal{P}_\alpha = \{\mathcal{U}_\alpha, \mathcal{X}_\alpha\}$ defined by the axiumbilic points \mathcal{U}_α and by the net \mathcal{X}_α (related to the crossing of principal axial curvature), in $M - \mathcal{U}_\alpha$ and the *mean axial configuration* $\mathcal{Q}_\alpha = \{\mathcal{U}_\alpha, \mathcal{Y}_\alpha\}$, defined by the axiumbilic points \mathcal{U}_α and the net \mathcal{Y}_α (related to the crossing of mean axial curvature), in $M - \mathcal{U}_\alpha$.




FIGURE 3. Projection on M of the integral curves of the Lie-Cartan vector field tangent to \mathbb{L}_α in a neighborhood of $\mathbf{p} \in M - \mathcal{U}_\alpha$. For each point in M pass four lines, associated, in pairs, to the axis of the ellipse.

2. DIFFERENTIAL EQUATION OF AXIAL LINES IN A MONGE CHART

The surface M will be locally parametrized by a Monge chart near an axiumbilic point \mathbf{p} as follows

$$z = R(x, y), \quad \text{and} \quad w = S(x, y),$$

where

$$(9) \quad R(x, y) = \frac{r_{20}}{2} x^2 + r_{11} xy + \frac{r_{02}}{2} y^2 + \frac{r_{30}}{6} x^3 + \frac{r_{21}}{2} x^2 y + \frac{r_{12}}{2} xy^2 + \frac{r_{03}}{6} y^3 \\ + \frac{r_{40}}{24} x^4 + \frac{r_{31}}{6} x^3 y + \frac{r_{22}}{4} x^2 y^2 + \frac{r_{13}}{6} xy^3 + \frac{r_{04}}{24} y^4 + h.o.t.,$$

$$(10) \quad S(x, y) = \frac{s_{20}}{2} x^2 + s_{11} xy + \frac{s_{02}}{2} y^2 + \frac{s_{30}}{6} x^3 + \frac{s_{21}}{2} x^2 y + \frac{s_{12}}{2} xy^2 + \frac{s_{03}}{6} y^3 \\ + \frac{s_{40}}{24} x^4 + \frac{s_{31}}{6} x^3 y + \frac{s_{22}}{4} x^2 y^2 + \frac{s_{13}}{6} xy^3 + \frac{s_{04}}{24} y^4 + h.o.t.$$

At the point $(x, y, R(x, y), S(x, y))$, the tangent plane to the surface is generated by $\{t_1, t_2\}$, where $t_1 = (1, 0, R_x, S_x)$ and $t_2 = (0, 1, R_y, S_y)$. The normal plane is generated by $\{N_1, N_2\}$, where $N_1 = \frac{\widetilde{N}_1}{|\widetilde{N}_1|}$ and $N_2 = \frac{\widetilde{N}_2}{|\widetilde{N}_2|}$ are defined by $\widetilde{N}_1 = (-R_x, -R_y, 1, 0)$ and $\widetilde{N}_2 = t_1 \wedge t_2 \wedge \widetilde{N}_1$.

Here \wedge is the *exterior* or *wedge* product $v_1 \wedge v_2 \wedge v_3$ of three vectors v_1, v_2, v_3 in \mathbb{R}^4 is defined by equation $\det(v_1, v_2, v_3, v) = \langle v_1 \wedge v_2 \wedge v_3, v \rangle$ for all $v \in \mathbb{R}^4$.

Therefore it follows that:

$$\det(t_1, t_2, \widetilde{N}_1, \bullet) = \langle \widetilde{N}_2, \bullet \rangle.$$

From the expressions of R and S given by equations (9) and (10), it follows that:

$$E = 1 + O(2), \quad F = O(2), \quad G = 1 + O(2),$$

and

$$\begin{aligned} e_1 &= r_{20} + r_{30}x + r_{21}y + O(2), & e_2 &= s_{20} + s_{30}x + s_{21}y + O(2), \\ f_1 &= r_{11} + r_{21}x + r_{12}y + O(2), & f_2 &= s_{11} + s_{21}x + s_{12}y + O(2), \\ g_1 &= r_{02} + r_{12}x + r_{03}y + O(2), & g_2 &= s_{02} + s_{12}x + s_{03}y + O(2). \end{aligned}$$

The axiumbilic points are defined by $a_0(x, y) = 0$ and $a_1(x, y) = 0$. So, in a neighborhood of $(0, 0)$, it follows that

$$(11) \quad a_0(x, y) = a_{00}^0 + a_{10}^0x + a_{01}^0y + O(2)$$

and

$$(12) \quad a_1(x, y) = a_{00}^1 + a_{10}^1x + a_{01}^1y + O(2),$$

where

$$\begin{aligned} a_{00}^0 &= r_{11}(r_{02} - r_{20}) + s_{11}(s_{02} - s_{20}), \\ a_{10}^0 &= r_{21}(r_{02} - r_{20}) + r_{11}(r_{12} - r_{30}) + s_{11}(s_{12} - s_{30}) + s_{21}(s_{02} - s_{20}), \\ a_{01}^0 &= r_{12}(r_{02} - r_{20}) + r_{11}(r_{03} - r_{21}) + s_{11}(s_{03} - s_{21}) + s_{12}(s_{02} - s_{20}) \end{aligned}$$

and

$$\begin{aligned} a_{00}^1 &= (r_{02} - r_{20})^2 + (s_{02} - s_{20})^2 - 4(r_{11}^2 + s_{11}^2), \\ a_{10}^1 &= 2(r_{12} - r_{30})(r_{02} - r_{20}) + 2(s_{12} - s_{30})(s_{02} - s_{20}) - 8(r_{21}r_{11} + s_{21}s_{11}), \\ a_{01}^1 &= 2(r_{03} - r_{21})(r_{02} - r_{20}) + 2(s_{03} - s_{21})(s_{02} - s_{20}) - 8(r_{12}r_{11} + s_{12}s_{11}). \end{aligned}$$

Therefore a point \mathbf{p} , expressed in a Monge chart by $(0, 0)$, is an axiumbilic point when the following relations hold.

$$(13) \quad \begin{cases} a_{00}^0 = r_{11}(r_{02} - r_{20}) + s_{11}(s_{02} - s_{20}) = 0, \\ a_{00}^1 = (r_{02} - r_{20})^2 + (s_{02} - s_{20})^2 - 4(r_{11}^2 + s_{11}^2) = 0. \end{cases}$$

Algebraic manipulations of the equations above, see [2], show that $(0, 0)$ is an axiumbilic point when the following equations hold

$$(14) \quad \begin{cases} 2r_{11} = (s_{02} - s_{20}), \\ 2s_{11} = -(r_{02} - r_{20}), \end{cases} \quad \text{or} \quad \begin{cases} 2r_{11} = -(s_{02} - s_{20}), \\ 2s_{11} = (r_{02} - r_{20}). \end{cases}$$

Remark 3. Let $r_{02} = r_{20} + r$ and $s_{02} = s_{20} + s$, $\rho^2 = r_{11}^2 + s_{11}^2$. Then condition for $(0, 0)$ to be an axiumbilic point, see equation (13), is given by

$$(15) \quad \begin{cases} r_{11} \cdot r + s_{11} \cdot s = 0, \\ r^2 + s^2 = 4\rho^2. \end{cases}$$

These condition for being an axiumbilic point can be interpreted as the intersection of a circle and a straight line in the plane (r, s) . The intersections are given by

$$(16) \quad \begin{cases} r_{11} = \frac{s}{2}, \\ s_{11} = -\frac{r}{2}, \end{cases} \quad \text{or} \quad \begin{cases} r_{11} = -\frac{s}{2}, \\ s_{11} = \frac{r}{2}, \end{cases}$$

and therefore equation (16) is another form of equation (14).

Let

$$\begin{aligned} \alpha_1 &= s_{12} - s_{30} + 2r_{21}, & \alpha_2 &= r_{30} - r_{12} + 2s_{21}, \\ \alpha_3 &= s_{03} - s_{21} + 2r_{12}, & \alpha_4 &= r_{21} - r_{03} + 2s_{12}. \end{aligned}$$

The discussion above is synthesized in the following lemma.

Lemma 4. Let \mathfrak{p} be an axiumbilic point with coordinates $(0, 0)$ in a Monge chart. The differential equation of axial lines in a neighborhood of $(0, 0)$ is given by

$$(17) \quad \tilde{a}_0(x, y)(dx^4 - 6dx^2dy^2 + dy^4) + \tilde{a}_1(x, y)(dx^2 - dy^2)dxdy + H(x, y, dx, dy) = 0,$$

where

$$(18) \quad \begin{aligned} \tilde{a}_0(x, y) &= \frac{1}{2}(r\alpha_1 + s\alpha_2)x + \frac{1}{2}(r\alpha_3 + s\alpha_4)y + a_{20}^0x^2 + a_{11}^0xy + a_{02}^0y^2, \\ \tilde{a}_1(x, y) &= 2(s\alpha_1 - r\alpha_2)x + 2(s\alpha_3 - r\alpha_4)y + a_{20}^1x^2 + a_{11}^1xy + a_{02}^1y^2 \end{aligned}$$

and H contains terms of order greater than or equal to 3 in (x, y) .

With the notation in equation (17), the *condition of transversality* between the curves $a_0 = 0$ and $a_1 = 0$ is given by

$$\begin{vmatrix} a_{10}^0 & a_{01}^0 \\ a_{10}^1 & a_{01}^1 \end{vmatrix} \neq 0.$$

The determinant above has the following expression:

$$[\alpha_2\alpha_3 - \alpha_1\alpha_4] \cdot (r^2 + s^2),$$

where $r = r_{02} - r_{20}$ and $s = s_{02} - s_{20}$. If $(r^2 + s^2)$ is zero, it follows that $a_{10}^0 = a_{01}^0 = a_{10}^1 = a_{01}^1 = 0$, and therefore the matrix

$$\begin{bmatrix} a_{10}^0 & a_{01}^0 \\ a_{10}^1 & a_{01}^1 \end{bmatrix}$$

is identically zero. Thus the axiumbilic points with $r = s = 0$ form a set of codimension at least four.

Therefore, the *condition of transversality*, supposing $r^2 + s^2 \neq 0$, is given by:

$$(19) \quad T := \alpha_2\alpha_3 - \alpha_1\alpha_4 \neq 0.$$

Long, but straightforward calculations show that condition (19) is invariant by positive rotations in the tangent and in the normal planes.

Lemma 5. Consider the quartic differential equation

$$(a_{10}x + a_{01}y)(dx^4 - 6dx^2dy^2 + dy^4) + (b_{10}x + b_{01}y)dxdy(dx^2 - dy^2) = 0.$$

Consider a rotation $x = \cos \theta u + \sin \theta v$, $y = -\sin \theta u + \cos \theta v$, where θ is a real root of the system of equations

$$-a_{01}t^5 + (a_{10} - b_{01})t^4 + (6a_{01} + b_{10})t^3 + (b_{01} - 6a_{10})t^2 - (a_{01} + b_{10})t + a_{10} = 0, \quad t = \tan \theta.$$

Then it follows that

$$\bar{a}_{01}v(du^4 - 6du^2dv^2 + dv^4) + (\bar{b}_{10}u + \bar{b}_{01}v)dudv(du^2 - dv^2) = 0, \text{ where}$$

$$\begin{aligned} a_{\bar{0}1} &= (t^2 + 1)[a_{01}(t^4 - 6t^2 + 1) + b_{01}t(t^2 - 1)] \\ b_{\bar{1}0} &= -16t(t^2 - 1)(a_{10} - a_{01}t) + (t^4 - 6t^2 + 1)(b_{10} - b_{01}t) \\ b_{\bar{0}1} &= -16(t^2 - 1)(ta_{10} + a_{01}) + (t^4 - 6t^2 + 1)(b_{10}t + b_{01}) \end{aligned}$$

Proof. The result follows from straightforward calculations. Observe that when $a_{01} = 0$ a rotation of angle $\pi/2$ is sufficient to obtain the result stated. \square

Proposition 6. Let \mathbf{p} be an axiumbilic point. Then there exists a Monge chart and a homothety in \mathbb{R}^4 such that the differential equation of axial lines is given by

$$(20) \quad y(dy^4 - 6dx^2dy^2 + dx^4) + (ax + by)dxdy(dx^2 - dy^2) + H(x, y, dx, dy) = 0$$

where H contains terms of order greater than or equal to 2 in (x, y) . Moreover, the axiumbilic point \mathbf{p} is transversal if and only if $a \neq 0$.

Proof. Consider a parametrization $X(x, y) = (x, y, R(x, y), S(x, y))$ given by equations (9) and (10) such that 0 is an axiumbilic point. By equation (18) it follows that:

$$\begin{aligned} a_0(x, y) &= \frac{1}{2}(r\alpha_1 + s\alpha_2)x + \frac{1}{2}(r\alpha_3 + s\alpha_4)y + O(2), \\ a_1(x, y) &= 2(s\alpha_1 - r\alpha_2)x + 2(s\alpha_3 - r\alpha_4)y + O(2). \end{aligned}$$

By an appropriate choice of the rotation in the plane $\{x, y\}$ given by Lemma 5 and a homothety in \mathbb{R}^4 , it is possible to make $2a_{10} = r\alpha_1 + s\alpha_2 = 0$ and, when $(\alpha_1\alpha_4 - \alpha_2\alpha_3)(r^2 + s^2) \neq 0$, also $a_{01} = \frac{1}{2}(r\alpha_3 + s\alpha_4) = 1$. So the result is established, $a = \frac{4(s\alpha_1 - r\alpha_2)}{r\alpha_3 + s\alpha_4}$ when $r\alpha_1 + s\alpha_2 = 0$ and $b = \frac{4(s\alpha_3 - r\alpha_4)}{r\alpha_3 + s\alpha_4}$. If $r \neq 0$ it follows that $a = -\frac{4(r^2 + s^2)\alpha_2}{r(r\alpha_3 + s\alpha_4)}$ and $a = \frac{4\alpha_1}{\alpha_4}$ when $s \neq 0$ and $r = 0$. \square

Remark 7. Let $p = \frac{dy}{dx}$. Then the differential equation (20) is given by:

$$(21) \quad y(p^4 - 6p^2 + 1) + (ax + by)p(1 - p^2) + H(x, y, p) = 0,$$

where H contains terms of order greater than or or equal to 2 in (x, y) .

3. AXIAL CONFIGURATION IN THE NEIGHBORHOOD OF AXIUMBILIC POINTS

Let \mathbf{p} be an axiumbilic point whose neighborhood is parametrized by a Monge chart and assume the notation established at the beginning of Section 2.

When it is a transversal axiumbilic point, which is determined by transversal intersection of the curves $a_0 = 0$ and $a_1 = 0$ (see equation (3)), it results from Proposition 6 and Remark 7 that the differential equation of axial lines is given by

$$(22) \quad \mathcal{G}(x, y, p) = y(p^4 - 6p^2 + 1) + (ax + by)p(1 - p^2) + H(x, y, p) = 0,$$

where $H(x, y, p)$ contains higher order terms greater or equal to 2 in (x, y) .

The Lie-Cartan surface \mathbb{L}_α in PM is defined implicitly by

$$(23) \quad \mathcal{G}(x, y, p) = 0.$$

In the case that \mathbf{p} is a transversal axiumbilic point the surface defined above is regular and of class \mathcal{C}^{r-2} in the neighborhood of the projective axis p .

In the coordinates (x, y, p) , the Lie-Cartan vector field X , is of class \mathcal{C}^{r-3} , (equation (8)):

$$(24) \quad X := \mathcal{G}_p \frac{\partial}{\partial x} + p\mathcal{G}_p \frac{\partial}{\partial y} - (\mathcal{G}_x + p\mathcal{G}_y) \frac{\partial}{\partial p}$$

and the projections of the integral curves of $X \Big|_{\mathcal{G}=0}$ are the axial lines in a neighborhood of \mathfrak{p} (Figure 3).

Restricted to the projective axis p , defined by $x = 0, y = 0$, the Lie-Cartan vector field is given by

$$X = -p[(p^4 - 6p^2 + 1) + (1 - p^2)(a + bp)] \frac{\partial}{\partial p}.$$

Therefore, the singular points of the Lie-Cartan vector field in the projective line are given by the equation:

$$(25) \quad P(p) = pR(p) = p[(p^4 - 6p^2 + 1) + (1 - p^2)(a + bp)] = 0.$$

The discriminant of $R(p) = (p^4 - 6p^2 + 1) + (1 - p^2)(a + bp)$ is

$$(26) \quad \begin{aligned} \Delta(a, b) = & 16a^5 + 4(b^2 + 68)a^4 + 16(b^2 + 144)a^3 \\ & - 8(b^2 - 80)(16 + b^2)a^2 + 96(16 + b^2)^2a + 4(16 + b^2)^3. \end{aligned}$$

Furthermore, $R(\pm 1) = -4$, $R(0) = 1 + a$ and $\lim_{p \rightarrow \pm\infty} R(p) = +\infty$, thus R has at least two simple real roots, one is less than -1 and the other is greater than 1 .

The derivative of X at $(0, 0, p)$ is given by:

$$DX(0, 0, p) = \begin{bmatrix} a(1 - 3p^2) & 4p^3 + b(1 - 3p^2) - 12p & 0 \\ a(1 - 3p^2)p & p[4p^3 + b(1 - 3p^2) - 12p] & 0 \\ 0 & 0 & -P'(p) \end{bmatrix}$$

whose eigenvalues are 0 and

$$\begin{aligned} \lambda_1(p) &= a(1 - 3p^2) + p[4p^3 + b(1 - 3p^2) - 12p], \\ \lambda_2(p) &= -P'(p). \end{aligned}$$

Recall that $P(p) = pR(p)$, and so $P'(p) = R(p) + pR'(p)$. Therefore at the roots of R , it follows that $-P'(p) = -pR'(p)$. Also, as ± 1 are not roots of R , it follows that

$$a = \frac{(-p^4 + 6p^2 - 1) + bp(1 - p^2)}{1 - p^2}.$$

Substituting the equation above into the expression of $\lambda_1(p)$, p being a root of $R(p)$ (a singular point of X), it follows that

$$\begin{cases} \lambda_1(p) = \frac{(p^2+1)^3}{(p^2-1)}, \\ \lambda_2(p) = -pR'(p). \end{cases}$$

Therefore, the eigenvalues of DX , at the singular points $(0, 0, p_0) = (0, 0, 0)$ and $(0, 0, p_i)$, $p_i \neq 0$, on the tangent space to $\mathcal{G} = 0$, are as follows:

$$(27) \quad p_0 = 0 : \begin{cases} \lambda_1 = a, \\ \lambda_2 = -(a + 1), \end{cases}$$

$$(28) \quad p_i \neq 0 : \begin{cases} \lambda_1 = \frac{(p_i^2+1)^3}{(p_i^2-1)}, \\ \lambda_2 = -p_i R'(p_i). \end{cases}$$

The eigenspace associated to the eigenvalue λ_1 is transversal to the axis p and the eigenvalue λ_2 has the projective axis as the associated eigenspace.

In [11] the axial configuration near an axiumbilic point was established in the following situation:

- $\Delta(a, b) < 0$,
- $\Delta(a, b) > 0$, $a < 0$, $a \neq -1$,
- $\Delta(a, b) > 0$, $a > 0$.

When $\Delta(a, b) < 0$, R has two simple real roots, and the Lie-Cartan vector field has three hyperbolic saddles in the projective axis. This axiumbilic point is called of type E_3 .

When $\Delta(a, b) > 0$, $a < 0$, $a \neq -1$, R has four simple real roots, and the Lie-Cartan vector field has 5 singular points in the projective line. Four are hyperbolic saddles and one is a hyperbolic node. This axiumbilic point is called of type E_4 .

When $\Delta(a, b) > 0$, $a > 0$, the Lie-Cartan vector field has 5 hyperbolic saddles in the projective line. This axiumbilic point is called of type E_5 .

In Figure 4 the Lie-Cartan surfaces and the integral curves of the Lie-Cartan vector field are sketched in the three cases E_3 , E_4 and E_5 . The projections of the integral curves by $\pi : PM \rightarrow M$ are the axial lines near the axiumbilic points (see Figure 5) E_3 , E_4 and E_5 .




FIGURE 4. Lie-Cartan vector field and its integral curves in the cases E_3 , E_4 and E_5 .




FIGURE 5. Axial configurations near axiumbilic points E_3 (left), E_4 (center) and E_5 (right).

For an immersion α of a surface M into \mathbb{R}^4 , the axiumbilic singularities \mathcal{U}_α and the lines of axial curvature are assembled into two *axial configurations*: the *principal axial configuration* $\mathcal{P}_\alpha = \{\mathcal{U}_\alpha, \mathcal{X}_\alpha\}$ and the *mean axial configuration* $\mathcal{Q}_\alpha = \{\mathcal{U}_\alpha, \mathcal{Y}_\alpha\}$.

An immersion $\alpha \in \mathcal{I}^r$ is said to be *principal axial stable* if it has a C^r neighborhood $\mathcal{V}(\alpha)$ such that, for any $\beta \in \mathcal{V}(\alpha)$ there exists a homeomorphism $h : M \rightarrow M$ mapping \mathcal{U}_α onto \mathcal{U}_β and mapping the integral net of \mathcal{X}_α onto that of \mathcal{X}_β . Analogous definition is given for *mean axial stability*.

In Proposition 8 are described the axiumbilic points which are principal axial stable. In Figure 6 are sketched the curves $\Delta(a, b) = 0$, $a = -1$ and $a = 0$ in the plane a, b , which bound the open regions corresponding to the three types of axiumbilic points of principal axial stable type.

Proposition 8 ([11], [12] p. 209). Let \mathbf{p} be an axiumbilic point of $\alpha \in \mathcal{I}^r$, $r \geq 5$. Then, α is locally principal axial stable and locally mean axial stable at \mathbf{p} if and only if \mathbf{p} is of type E_3 , E_4 or E_5 . The curve $\Delta(a, b) = 0$ has three connected components, is contained in the region $a \leq -1$ and it is regular outside the points $(-\frac{27}{2}, \pm \frac{5\sqrt{5}}{2})$ which are of cuspidal type.




FIGURE 6. Diagram of stable axiumbilic points, E_3 , E_4 and E_5 .

Proof. The function $\Delta(a, b)$ defined by equation (26) is symmetric in b . The polynomials $\Delta(a, b)$ and $\frac{\partial \Delta}{\partial b}$ in the variable b have resultant equal to a positive multiple of

$$(1 + a)(a^2 + 8a + 32)^2 a^{16} (2a + 27)^6.$$

The critical points $p_\pm = (-\frac{27}{2}, \pm \frac{5\sqrt{5}}{2})$ of Δ are contained in $\Delta(a, b) = 0$. Near the point p_+ it follows that:

$$\Delta(a, b) = -54675 \left[\left(a + \frac{27}{2} \right)^2 + 5 \left(b - \frac{5\sqrt{5}}{2} \right)^2 + 2\sqrt{5} \left(a + \frac{27}{2} \right) \left(b - \frac{5\sqrt{5}}{2} \right) \right] + h.o.t.$$

Further analysis shows p_\pm are Whitney cuspidal points.

Also the curve $\Delta(a, b) = 0$ is contained in the region $a \leq -1$ and near $(-1, 0)$ it is given by $a = -\frac{1}{20}b^2 + O(3)$. In fact, for $a > -1$ all the roots of $\Delta(a, b)$ are complex.

By the classification of axiumbilic points E_3 , E_4 and E_5 by the sign of $\Delta(a, b)$ and of a , the diagram of stable axiumbilic points, see [11], [12] p. 209, is as shown in Fig. 6. \square

3.1. The axiumbilic point E_{34}^1 .

Definition 9. Let $\alpha : M \rightarrow \mathbb{R}^4$ be an immersion of class C^r , $r \geq 5$, of a smooth and oriented surface. An axiumbilic point \mathbf{p} is said to be of type E_{34}^1 if a defined in Proposition 6 does not vanish and:

- i) $\Delta(a, b) = 0$, $(a, b) \neq (-1, 0)$ and $(a, b) \neq (-\frac{27}{2}, \pm\frac{5}{2}\sqrt{5})$, or
- ii) $b \neq 0$ if $a = -1$.

Proposition 10. Let $\alpha : M \rightarrow \mathbb{R}^4$ be an immersion of class C^r , $r \geq 5$ of a smooth and oriented surface having an axiumbilic point \mathbf{p} of type E_{34}^1 . Then the axial configuration, defined in subsection 1.1, of α in a neighborhood of \mathbf{p} is as shown in Figure 7.




FIGURE 7. Axial configurations in a neighborhood of an axiumbilic point of type E_{34}^1 .

Proof. Since the condition of transversality ($a \neq 0$) is preserved at an axiumbilic point of type E_{34}^1 the implicit surface defined by equation (23) is regular in a neighborhood of the projective line. From the hypotheses $\Delta(a, b) = 0$, $(a, b) \neq (-1, 0)$ and $(a, b) \neq (-\frac{27}{2}, \pm\frac{5}{2}\sqrt{5})$ or $b \neq 0$, if $a = -1$, the polynomial $P(p) = p[(p^4 - 6p^2 + 1) + (1 - p^2)(a + bp)] = pR(p)$, which defines the singularities of the Lie-Cartan vector field, has one double root and three real simple roots.

With no loss of generality, we can consider the case $a = -1$ and $b \neq 0$, where $p = 0$ is a double root of the polynomial $P(p)$. In this case, we have $P(p) = p^2(p^3 - bp^2 - 5p + b)$. The eigenvalues of DX at $(0, 0, p)$ are given by:

$$\lambda_1 = 4p^4 - 3bp^3 - 9p^2 + bp - 1 \quad \text{and} \quad \lambda_2 = p(-5p^3 + 4bp^2 + 15p - 2b).$$

Therefore, at the singular points $(0, 0, p)$, $p \neq 0$, of X it follows that:

$$\lambda_1 = \frac{(p^2 + 1)^3}{p^2 - 1} \quad \text{and} \quad \lambda_2 = -\frac{p^2(p^4 + 2p^2 + 5)}{p^2 - 1}.$$

Then, $\lambda_1 \lambda_2 < 0$ when $p \neq 0$ and these three singular points of X are hyperbolic saddles. At $p = 0$, double root of P , it follows that $\lambda_1 = -1$, $\lambda_2 = 0$. Recall that the eigenspace associated to λ_1 is transversal to the axis p and that one associated to λ_2 is the projective axis itself. Since $\mathcal{G}_y(0, 0, 0) = 1$, it follows from the implicit function theorem that $y(x, p) = xp + O(3)$ is defined in a neighborhood of $(0, 0, 0)$ such that $\mathcal{G}(x, y(x, p), p) = 0$. In this case, the Lie-Cartan vector field in the chart (x, p) is given by:

$$(29) \quad \begin{cases} \dot{x} = -x + bxp + O(3) \\ \dot{p} = -bp^2 + O(3) \end{cases}$$

with $b \neq 0$. Therefore, $(0, 0, 0)$ is a quadratic saddle-node with the center manifold tangent to the projective line. The phase portrait is sketched in Figure 8, and the projections of the integral curves are the axial lines shown in Figure 7.




FIGURE 8. Integral curves of $X|_{G=0}$ in the neighborhood of the projective line in the case of an axiumbilic point of type E_{34}^1

When $(a, b) \neq (-1, 0)$, $(a, b) \neq (-\frac{27}{2}, \pm\frac{5}{2}\sqrt{5})$ and $\Delta(a, b) = 0$ the polynomial

$$P(p) = p[(p^4 - 6p^2 + 1) + (1 - p^2)(a + bp)]$$

has a double root $p_0 \neq 0$ and three real simple roots. This case is reduced to the case when $p = 0$ is a double root, making an appropriate rotation of coordinates in the plane $\{x, y\}$ so that, in the new coordinates, the double root p_0 is located at $p = 0$. \square

Proposition 11. Let $\alpha \in \mathcal{I}^r$, $r \geq 5$, be an immersion such that \mathfrak{p} is axiumbilic point of type E_{34}^1 . Then, there is a neighborhood V of \mathfrak{p} , a neighborhood \mathcal{V} of α and a function $\mathcal{F} : \mathcal{V} \rightarrow \mathbb{R}$ of class \mathcal{C}^{r-3} such that for each $\mu \in \mathcal{V}$ there is an unique axiumbilic point $\mathfrak{p}_\mu \in V$ such that:

- i) $d\mathcal{F}_\alpha \neq 0$,
- ii) $\mathcal{F}(\mu) < 0$ if and only if \mathfrak{p}_μ is of type E_3 ,
- iii) $\mathcal{F}(\mu) > 0$ if and only if \mathfrak{p}_μ of type E_4 ,
- iv) $\mathcal{F}(\mu) = 0$ if, and only if, \mathfrak{p}_μ is of type E_{34}^1 .

Proof. Since \mathfrak{p} is a transversal axiumbilic point of α , the existence of the neighborhoods \mathcal{V} and V follows from the Implicit Function Theorem. For $\mu \in \mathcal{V}$ with an axiumbilic point $\mathfrak{p}_\mu \in V$, after a rigid motion Γ_μ in \mathbb{R}^4 , locally the immersion $\mu \in \mathcal{V}$ can be parametrized in terms of a Monge chart $(x, y, R_\mu(x, y), S_\mu(x, y))$, with the origin being the axiumbilic point p_μ and

$$\begin{aligned} R_\mu(x, y) &= \frac{r_{20}(\mu)}{2}x^2 + r_{11}(\mu)xy + \frac{r_{02}(\mu)}{2}y^2 + \frac{r_{30}(\mu)}{6}x^3 + \frac{r_{31}(\mu)}{2}x^2y \\ &\quad + \frac{r_{13}(\mu)}{2}xy^2 + \frac{r_{03}(\mu)}{6}y^3 + h.o.t., \\ S_\mu(x, y) &= \frac{s_{20}(\mu)}{2}x^2 + s_{11}(\mu)xy + \frac{s_{02}(\mu)}{2}y^2 + \frac{s_{03}(\mu)}{6}x^3 + \frac{s_{21}(\mu)}{2}x^2y \\ &\quad + \frac{s_{12}(\mu)}{2}xy^2 + \frac{s_{03}(\mu)}{6}y^3 + h.o.t. \end{aligned}$$

For μ , performing rotations and homoteties as described in Section 2, the coefficients a_μ and b_μ can be expressed in function of the coefficients of the surface presented in a Monge chart, as was done in Proposition 6, considering the coefficients in function of the parameter $\mu \in \mathcal{V}$.

Define $\mathcal{F}(\mu) = \Delta(a(\mu), b(\mu))$ whose zeros define locally the manifold of immersions with an E_{34}^1 axiumbilic point. Here, $\Delta(a, b)$, given by equation (26), is the discriminant of the polynomial $R(p) = (p^4 - 6p^2 + 1) + (1 - p^2)(a + bp)$.

Notice that due to the particular representation of the 3-jets taken here, the condition $a(\mu) = -1$ in Definition 9, the jet extension of the immersion is not transversal, but tangent, to

the manifold of jets with E_{34}^1 axiumbilic points. It is always possible, by an appropriate rotation in the plane $\{x, y\}$ to suppose that $a(\alpha) \notin \{-\frac{27}{2}, -1\}$. See Section 2.

Assertions (ii), (iii) and (iv) follow from the definition of \mathcal{F} and the previous analysis on the sign of the discriminant $\Delta(a_\mu, b_\mu)$.

Moreover, the derivative of $\mathcal{F}(\mu)$ in the direction of the coordinate a does not vanish, leading to conclude that $d\mathcal{F}_\alpha \neq 0$.

In fact, assuming $s_{11}(\alpha) = \frac{1}{2}r \neq 0$, it follows that $a_0(\mu) = y + O(2)$,

$$\begin{aligned} a_1(\mu)(x, y) &= -\frac{4(r(\mu)^2 + s(\mu)^2)\alpha_2(\mu)}{r(\mu)(r(\mu)\alpha_3(\mu) + s(\mu)\alpha_4(\mu))}x + \frac{4(s(\mu)\alpha_3(\mu) - r(\mu)\alpha_4(\mu))}{r(\mu)\alpha_3(\mu) + s(\mu)\alpha_4(\mu)}y + O(2) \\ &= a(\mu)x + b(\mu)y + O(2), \end{aligned}$$

$\alpha_1 = s_{12} - s_{30} + 2r_{21}$, $\alpha_2 = r_{30} - r_{12} + 2s_{21}$, $\alpha_3 = s_{03} - s_{21} + 2r_{12}$, and $\alpha_4 = r_{21} - r_{03} + 2s_{12}$.

Consider the deformation

$$\mu = (x, y, R_\alpha(x, y), S_\alpha(x, y)) + \left(0, 0, t\left(\frac{1}{6}x^3 - \frac{1}{2}xy^2\right), tx^2y\right).$$

Then, as $\alpha_2 = r_{30} - r_{12} + 2s_{21}$, it follows that $a(\mu) = -\frac{4(r^2+s^2)(\alpha_2+t)}{r(r\alpha_3+s\alpha_4)}$ and

$$\left.\frac{d}{dt}(\Delta(a(\mu), b(\mu)))\right|_{t=0} = \frac{\partial\Delta}{\partial a} \cdot \frac{da}{dt} = \frac{\partial\Delta}{\partial a} \cdot \left(-\frac{4(r^2+s^2)}{r(r\alpha_3+s\alpha_4)}\right) \neq 0.$$

In the case where $s_{11}(\alpha) = 0$ it follows that $r_{11}(\alpha) = -\frac{1}{2}s \neq 0$, $\alpha_1\alpha_4 \neq 0$ and $\alpha_2(\mu) = 0$. Now consider the deformation

$$\mu = (x, y, R_\alpha(x, y), S_\alpha(x, y)) + \left(0, 0, tx^2y, t\left(-\frac{1}{6}x^3 + \frac{1}{2}xy^2\right)\right).$$

Then, $a(\mu) = \frac{4(\alpha_1+t)}{\alpha_4}$ and

$$\left.\frac{d}{dt}(\Delta(a(\mu), b(\mu)))\right|_{t=0} = \frac{\partial\Delta}{\partial a} \cdot \frac{da}{dt} = \frac{\partial\Delta}{\partial a} \cdot \left(\frac{4}{\alpha_4}\right) \neq 0.$$

□




FIGURE 9. Axial configuration near axiumbilic points. E_3 (left), E_{34}^1 (center) and E_4 (right).




FIGURE 10. Bifurcation diagram of the axial configuration near an axiumbilic point E_{34}^1 and the structure of separatrices.

3.2. **The axiumbilic point $E_{4,5}^1$.** Consider the Monge chart described by equations (9) and (10). Suppose that the origin is an axiumbilic point, which is expressed by

$$(30) \quad \begin{aligned} R(x, y) = & \frac{r_{20}}{2}x^2 + r_{11}xy + \frac{r_{02}}{2}y^2 + \frac{r_{30}}{6}x^3 + \frac{r_{21}}{2}x^2y + \frac{r_{12}}{2}xy^2 + \frac{r_{03}}{6}y^3 \\ & + \frac{r_{40}}{24}x^4 + \frac{r_{31}}{6}x^3y + \frac{r_{22}}{4}x^2y^2 + \frac{r_{13}}{6}xy^3 + \frac{r_{04}}{24}y^4 + h.o.t., \end{aligned}$$

$$(31) \quad \begin{aligned} S(x, y) = & \frac{s_{20}}{2}x^2 + s_{11}xy + \frac{s_{02}}{2}y^2 + \frac{s_{30}}{6}x^3 + \frac{s_{21}}{2}x^2y + \frac{s_{12}}{2}xy^2 + \frac{s_{03}}{6}y^3 \\ & + \frac{s_{40}}{24}x^4 + \frac{s_{31}}{6}x^3y + \frac{s_{22}}{4}x^2y^2 + \frac{s_{13}}{6}xy^3 + \frac{s_{04}}{24}y^4 + h.o.t., \end{aligned}$$

where, $r_{02} = r_{20} + r$, $r_{11} = -\frac{1}{2}s$, $s_{02} = s_{20} + s$, $s_{11} = \frac{1}{2}r$.

Let $\alpha_1 = s_{12} - s_{30} + 2r_{21}$, $\alpha_2 = r_{30} - r_{12} + 2s_{21}$, $\alpha_3 = s_{03} - s_{21} + 2r_{12}$, $\alpha_4 = r_{21} - r_{03} + 2s_{12}$, $\beta_1 = s_{22} - s_{40} + 2r_{31}$, $\beta_2 = r_{40} - r_{22} + 2s_{31}$, $\beta_3 = s_{13} - s_{31} + 2r_{22}$, $\beta_4 = r_{31} - r_{13} + 2s_{22}$, $\beta_5 = s_{04} - s_{22} + 2r_{13}$, and $\beta_6 = r_{22} - r_{04} + 2s_{13}$.

The functions a_0 and a_1 (see Proposition 1) are given by

$$(32) \quad a_0(x, y) = a_{10}x + a_{01}y + \frac{1}{2}a_{20}x^2 + a_{11}xy + \frac{1}{2}a_{02}y^2 + h.o.t. \quad \text{and}$$

$$(33) \quad a_1(x, y) = b_{10}x + b_{01}y + \frac{1}{2}b_{20}x^2 + b_{11}xy + \frac{1}{2}b_{02}y^2 + h.o.t.,$$

where

$$\begin{aligned} a_{10} &= \frac{1}{2}(r\alpha_1 + s\alpha_2), & a_{01} &= \frac{1}{2}(r\alpha_3 + s\alpha_4), \\ a_{20} &= -\alpha_2r_{21} + \alpha_1s_{21} + \left[\frac{\beta_1}{4} + \frac{s_{20}}{2}(r_{20}^2 + s_{20}^2) \right] r + \\ & \quad \left[\frac{\beta_2}{4} - \frac{r_{20}}{2}(r_{20}^2 + s_{20}^2) \right] s + (r_{20}^2 - s_{20}^2)sr - \frac{3}{8}(r^2 + s^2)(s_{20}r - r_{20}s) + r_{20}s_{20}(s^2 - r^2), \\ a_{11} &= -r_{12}\alpha_2 + s_{12}\alpha_1 - r_{21}\alpha_4 + s_{21}\alpha_3 - \left[\frac{\beta_3}{2} + r_{20}(r_{20}^2 + s_{20}^2) \right] r + \left[\frac{\beta_4}{2} - s_{20}(r_{20}^2 + s_{20}^2) \right] s \\ & \quad - 2s_{20}r_{20}rs - \frac{1}{2}(3s_{20}^2 + r_{20}^2)s^2 - \frac{1}{2}(3r_{20}^2 + s_{20}^2)r^2 - \frac{3}{8}(r^2 + s^2)^2 - \frac{5}{4}(r^2 + s^2)(r_{20}r + s_{20}s), \end{aligned}$$

$$\begin{aligned}
a_{02} &= -r_{12}\alpha_4 + s_{12}\alpha_3 + \left[\frac{\beta_5}{2} - \frac{s_{20}}{2}(r_{20}^2 + s_{20}^2) \right] r + \left[\frac{\beta_6}{2} + \frac{r_{20}}{2}(r_{20}^2 + s_{20}^2) \right] s + \\
&\quad (-2s_{20}^2 + 2r_{20}^2)sr + 2s_{20}r_{20}(s^2 - 2r^2) + -\frac{9}{8}(r^2 + s^2)(rs_{20} - sr_{20}), \\
b_{10} &= 2(s\alpha_1 - r\alpha_2), \quad b_{01} = 2(s\alpha_3 - r\alpha_4), \\
b_{20} &= \alpha_1^2 + \alpha_2^2 - 4(s_{21}\alpha_2 + r_{21}\alpha_1) + \left[-\beta_2 + 2r_{20}(r_{20}^2 + s_{20}^2) \right] r + \\
&\quad \left[\beta_1 + 2s_{20}(r_{20}^2 + s_{20}^2) \right] s - \frac{1}{2}(r^2 + s^2)(s_{20}s + r_{20}r) + 4(r_{20}s - s_{20}r)^2, \\
b_{11} &= 2(\alpha_3\alpha_1 + \alpha_2\alpha_4) - 4(\alpha_1r_{12} + \alpha_2s_{12} + \alpha_3r_{21} + \alpha_4s_{21}) + \\
&\quad 2 \left[-\beta_4 + 2s_{20}(r_{20}^2 + s_{20}^2) \right] r + 2 \left[\beta_3 - 2r_{20}(r_{20}^2 + s_{20}^2) \right] s + 4(s_{20}^2 - r_{20}^2)rs + 4r_{20}s_{20}(r^2 - s^2), \\
b_{02} &= \alpha_3^2 + \alpha_4^2 + 4(r_{12}^2 + s_{12}^2) + 4s_{12}(r_{21} - r_{03}) + 4r_{12}(s_{03} - s_{21}) + \\
&\quad [-\beta_6 - 2r_{20}(r_{20}^2 + s_{20}^2)]r + [\beta_5 - 2s_{20}(r_{20}^2 + s_{20}^2)]s + 2(r_{20}^2 - 3s_{20}^2)s^2 + 2(s_{20}^2 - r_{20}^2)r^2.
\end{aligned}$$

Definition 12. An axiumbilic point is said to be of type $E_{4,5}^1$ if the variety \mathbb{L}_α has exactly 4 singular points which are of Morse type located along the projective line over the point.

Proposition 13. Consider a Monge chart and a homothety such that the differential equation of axial lines is written as

$$a_0(x, y)(dx^4 - 6dx^2dy^2 + dy^4) + a_1(x, y)dxdy(dx^2 - dy^2) + O(3) = 0,$$

where

$$\begin{aligned}
a_0(x, y) &= y + \frac{1}{2}a_{20}x^2 + a_{11}xy + \frac{1}{2}a_{02}y^2 + h.o.t., \\
a_1(x, y) &= b_{01}y + \frac{1}{2}b_{20}x^2 + b_{11}xy + \frac{1}{2}b_{02}y^2 + h.o.t.
\end{aligned}$$

Then the following conditions are equivalent:

- i) the curves $a_0 = 0$ and $a_1 = 0$ are regular and have quadratic contact at 0,
- ii) the axiumbilic point 0 is of type $E_{4,5}^1$,
- iii) the Lie-Cartan vector field defined in \mathbb{L}_α has a quadratic saddle-node in the projective axis with the center manifold transversal to the projective line.

Proof. The differential equation of axial lines can be written as

$$a_0(x, y)(dx^4 - 6dx^2dy^2 + dy^4) + a_1(x, y)dxdy(dx^2 - dy^2) + O(3) = 0,$$

where

$$\begin{aligned}
a_0(x, y) &= a_{10}x + a_{01}y + \frac{1}{2}a_{20}x^2 + a_{11}xy + \frac{1}{2}a_{02}y^2 + h.o.t. \\
a_1(x, y) &= b_{10}x + b_{01}y + \frac{1}{2}b_{20}x^2 + b_{11}xy + \frac{1}{2}b_{02}y^2 + h.o.t.
\end{aligned}$$

where the coefficients of a_0 and a_1 are given by equations (32) and (33). Here $O(3)$ means terms of order greater than or equal to 3 in the variables x and y .

In what follows it will be considered a Monge chart such that $a_{10} = 0$. This is possible as shown in Lemma 5 and Proposition 6. Since the contact between $a_0 = 0$ and $a_1 = 0$ is supposed

to be quadratic it results that $b_{10} = 0$ and $a_{01} \cdot b_{01} \neq 0$. Also by a homothety it is possible to obtain $a_{01} = 1$.

So, it results that:

$$(34) \quad a_0(x, y) = y + \frac{1}{2}a_{20}x^2 + a_{11}xy + \frac{1}{2}a_{02}y^2 + h.o.t.$$

$$(35) \quad a_1(x, y) = b_{01}y + \frac{1}{2}b_{20}x^2 + b_{11}xy + \frac{1}{2}b_{02}y^2 + h.o.t.$$

Therefore, the condition of quadratic contact between the two regular curves is expressed by $\chi = b_{20} - a_{20}b_{01} \neq 0$.

Notice that this amounts to establish the implication $i) \rightarrow ii)$.

Claim 14. In the neighborhood of $(0, 0, 0)$, the Lie-Cartan vector field restricted to the surface $\mathcal{G} = 0$, can be expressed in the chart (x, p) by

$$(36) \quad \begin{cases} \dot{x} = \frac{\chi}{2}x^2 + O(3), \\ \dot{p} = -p + \frac{3}{2}a_{11}a_{20}x^2 - (a_{11} + \chi)p - b_{01}p^2 + O(3) \end{cases}$$

and $(0, 0, 0)$ is a saddle-node when $\chi \neq 0$.

Since $\mathcal{G}_y(0, 0, 0) = 1 \neq 0$, it follows from implicit function theorem that locally $y = y(x, p)$ and $\mathcal{G}(x, y(x, p), p) = 0$.

The Taylor expansion of $y(x, p)$ in the neighborhood of $(x, p) = (0, 0)$ is given by:

$$(37) \quad y(x, p) = -\frac{1}{2}a_{20}x^2 + O(3).$$

The Lie-Cartan vector field restricted to the surface $\mathcal{G} = 0$ is given by

$$\begin{cases} \dot{x} = \mathcal{G}_p(x, y(x, p), p) = \frac{1}{2}\chi x^2 + O(3) \\ \dot{p} = -(\mathcal{G}_x + p\mathcal{G}_y)(x, y(x, p), p) = -p + \frac{3}{2}a_{11}a_{20}x^2 - (\chi + a_{11})p - b_{01}p^2 + O(3) \end{cases}$$

The eigenvalues of the vector field (36) at 0 are $\lambda_1 = 0$ and $\lambda_2 = -1$ with respective associated eigenspaces $\ell_1 = (1, -a_{20})$ and $\ell_2 = (0, 1)$. By Invariant Manifold Theory the center manifold is tangent to ℓ_1 and is given by $W^c = \{(x, -a_{20}x + \frac{3}{2}a_{20}(\chi + a_{11})x^2 + O(3))\}$.

The restriction of the vector field (36) to the center manifold is given by $[\frac{1}{2}\chi x^2 + O(3)]\frac{\partial}{\partial x}$.

This establishes that $ii) \rightarrow iii)$.

Claim 15. The function \mathcal{G} has exactly 4 critical points in the projective line, and they are of Morse-type of index 1 or 2 if and only if $\chi \neq 0$.

The critical points of \mathcal{G} along the projective line are determined by

$$(38) \quad S(p) = \mathcal{G}_v(0, 0, p) = (p^4 - 6p^2 + 1) + b_{01}p(1 - p^2) = 0,$$

which has for 4 simple real roots located in the intervals $(-\infty, -1)$, $(-1, 0)$, $(0, 1)$ and $(1, \infty)$. This follows from $S(\pm 1) = -4$, $S(0) = 1$ and from the discriminant $\Delta(S) = 4(16 + b_{01}^2)^3 > 0$.

Along the projective line, the determinant of the Hessian of \mathcal{G} is given by

$$(39) \quad \text{Hess } \mathcal{G}(0, 0, p) = -(a_{20}(1 - 6p^2 + p^4) + b_{20}p(1 - p^2))(b_{01} - 12p - 3b_{01}p^2 + 4p^3)^2.$$

The resultant of $S(p)$ and $\text{Hess } \mathcal{G}(0, 0, p)$ is given by $256\chi^4(16 + b_{01}^2)^6$ and therefore $\text{Hess } \mathcal{G}(0, 0, p) \neq 0$ at the critical points of \mathcal{G} . This implies that the critical points are of Morse type. As $\mathcal{G}(0, 0, p) = 0$ it follows that the index of a critical point is 1 or 2 and so locally the level set $\mathcal{G} = 0$ is a cone.

The eigenvalues of the derivative of the Lie-Cartan vector field at a point $(0, 0, p)$ are given by:

$$\lambda_1 = -p(-4p^3 + 3b_{01}p^2 + 12p - b_{01}), \quad \lambda_2 = -1 + 18p^2 - 5p^4 - 2b_{01}p + 4b_{01}p^3.$$

At the critical points p_i (satisfying $S(p_i) = 0$) it follows that $\lambda_1 = -\lambda_2 = \frac{p^6 + 3p^4 + 3p^2 + 1}{p^2 - 1}$, then $\lambda_1^i \lambda_2^i < 0$, for $i = 1..4$.

Therefore, these 4 points are saddles of the Lie-Cartan vector field. As the projective line is invariant it follows that the other invariant manifold (stable or stable) of a singular point is transversal to the projective line.

This amounts to prove that $iii) \rightarrow i)$. □

Proposition 16. Let $\alpha \in \mathcal{I}^r$, $r \geq 5$ and \mathbf{p} be an axiumbilic point. Suppose, in the Monge chart expressed by equations (30) and (31), that $\alpha_1 = \alpha_3 = 0$ and $\chi \neq 0$. Then \mathbf{p} is an axiumbilic point of type $E_{4,5}^1$ and the axial configurations of α in a neighborhood of \mathbf{p} is as shown in Figure 11.



FIGURE 11. Axial configurations in a neighborhood of an axiumbilic point of type $E_{4,5}^1$.




FIGURE 12. Lie-Cartan vector field near an axiumbilic point $E_{4,5}^1$ and the axial configuration (principal and mean).

Proof. Condition $\alpha_1 = \alpha_3 = 0$ implies the non-transversal contact of the curves $a_0 = 0$ and $a_1 = 0$ at the axiumbilic point \mathfrak{p} expressed in the Monge chart by $(0, 0)$. By Lemma 5 and Proposition 6, it is possible to express these curves as in equation (34). Assuming $\chi \neq 0$, we have the quadratic contact of the curves at the axiumbilic point.

Proposition 13 implies that over the axiumbilic point we have five equilibria of the Lie-Cartan vector field. One of them is a regular point of the Lie-Cartan surface, and this is an equilibrium of saddle-node type with center manifold transversal to the axis p (see Claim 14).

The remaining equilibria are critical points of Morse type of the Lie-Cartan surface. In the neighborhood of these points, the level set $\mathcal{G} = 0$ are locally cones, and the 4 points are saddles of the Lie-Cartan vector field (see Claim 15).

Therefore, we conclude that the configuration is as described in Figure 12, whose projection of the saddle-node and parallel sectors describe the principal axial and mean axial configurations close to the axiumbilic point \mathfrak{p} of type E_{45}^1 (Figure 11). \square

Proposition 17. Let $\alpha \in \mathcal{I}^r$, $r \geq 5$, be an immersion having an axiumbilic point \mathfrak{p} of type $E_{4,5}^1$. Then, there exist a neighborhood V of \mathfrak{p} , a neighborhood \mathcal{V} of α and a function $F : \mathcal{V} \rightarrow \mathbb{R}$ of class \mathcal{C}^{r-3} such that:

- i) $dF_\alpha \neq 0$,
- ii) $F(\mu) = 0$ if, and only if, $\mu \in \mathcal{V}$ has just one axiumbilic point in V , which is of type $E_{4,5}^1$,
- iii) $F(\mu) < 0$ if, and only if, μ has exactly two axiumbilic points in V , one of type E_4 and the other of type E_5 ,
- iv) $F(\mu) > 0$ if, and only if, μ has no axiumbilic points in V .

Proof. By Proposition 13, α being an immersion having an axiumbilic point \mathfrak{p} of type $E_{4,5}^1$, the curves $a_0^\alpha = 0$ and $a_1^\alpha = 0$ have quadratic contact at \mathfrak{p} .

Since $\frac{\partial a_0^\alpha}{\partial y}(0, 0) = a_{01} \neq 0$, it follows from Implicit Function Theorem that locally, for μ in a neighborhood \mathcal{V} of α , $y = y_\mu(x)$ and $a_0^\mu(x, y_\mu(x)) = 0$.

Moreover, $\frac{\partial^2 a_1^\alpha}{\partial x^2}(0, 0) = b_{20} \neq 0$, and so $x = x_\mu$ is a local solution of $\frac{\partial a_1^\mu}{\partial x}(x_\mu, y_\mu(x_\mu)) = 0$. Define $F(\mu) = a_1^\mu(x_\mu, y_\mu(x_\mu))$. Consider the variation

$$h_t(x, y) = (x, y, R(x, y) + txy, S(x, y) + txy).$$

It follows that $\left. \frac{dF(t)}{dt} \right|_{t=0} \neq 0$, and so $dF_\alpha \neq 0$. Therefore, the result follows from the Implicit Function Theorem.

The axiumbilic point of type $E_{4,5}^1$ is therefore the transition between zero and two axiumbilic points, one of type E_4 and the other of type E_5 .

In Figures 13 and 15 are illustrated this transition, with the axial configurations sketched in two different styles. See also Figure 15 for an illustration of transition in the Lie - Cartan surface. \square

Proposition 18. In the space of smooth mappings of $M \times \mathbb{R} \rightarrow \mathbb{R}^4$ which are immersions relative to the first variable, those which have all their axiumbilic points either generic (of types E_3 , E_4 and E_5) or of types E_{34}^1 and E_{45}^1 , crossed transversally, is open and dense. Furthermore, for such families the axiumbilic points describe a regular curve in $M \times \mathbb{R}$ whose projection into \mathbb{R} has only non-degenerate critical points at $E_{4,5}^1$ and the regular points of the projection is a collection of arcs bounded by E_{34}^1 points, which are the common boundary points of the arcs consisting of points of types E_3 and E_4 .

Proposition 18 follows from the analysis in Propositions 11 and 17 and an application of Thom Transversality Theorem to the submanifold of four jets of immersions at axiumbilic points,




FIGURE 13. Axiumbilic point E_{45}^1 . The axiumbilic points E_4 and E_5 collapse in an axiumbilic point E_{45}^1 , and after they are eliminated and there are no axiumbilic points.




FIGURE 14. Bifurcation diagram of the axial configuration near an axiumbilic point of type E_{45}^1 and the structure of separatrices




FIGURE 15. The Lie-Cartan surface. In the left, with two axiumbilic point, in the center with four singular points, and in the right the four regular levels.

stratified by the generic axiumbilic points of types E_3 , E_4 and E_5 , by those of types E_{34}^1 and E_{45}^1 , and by their complement which has codimension larger than 3. See Section 4.

4. TRANSVERSALITY AND STRATIFICATION

Consider the space $\mathbb{J}^k(M, \mathbb{R}^4)$ of k -jets of immersions α of a compact oriented surface M into \mathbb{R}^4 , endowed with the structure of Principal Fiber Bundle. The base is M ; the fiber is the space $\mathbb{R}^4 \times \mathbb{J}^k(2, 4)$, where $\mathcal{J}^k(2, 4)$ is the space of k -jets of immersions of \mathbb{R}^2 to \mathbb{R}^4 , preserving the respective origins. The structure group, \mathbb{A}_+^k , is the product of the group $\mathcal{L}_+^k(2, 2)$ of k -jets of origin and orientation preserving diffeomorphisms of \mathbb{R}^2 , acting on the right by coordinate changes, and by the group of positive isometries of \mathbb{R}^4 , acting on the left. This group is generated by the groups of translations and that of positive rotations, $\mathcal{O}_+(4)$, of \mathbb{R}^4 .

Denote by $\Pi_{k,l}, k \leq l$ the projection of $\mathcal{J}^l(2, 4)$ to $\mathcal{J}^k(2, 4)$. It is well known that the group action commutes with projections.

Definition 19. We define below the *canonic axiumbilic stratification* of $\mathcal{J}^4(2, 4)$. The term *canonic* means that the strata are invariant under the action of the group $\mathbb{A}_+^k = \mathcal{O}_+(4) \times \mathcal{L}_+^k(2, 2)$.

- 1) *Axiumbilic Jets:* \mathcal{U}^4 , those in the orbit of $j^4(x, y, R(x, y), S(x, y))$, where R and S are as in equations (9) and (10) satisfying the axiumbilic conditions defined in terms of $j^2R(0)$ and $j^2S(0)$. It is a closed variety of codimension 2.
- 2) *Non-axiumbilic Jets:* $(\mathcal{N}\mathcal{U})^4$ is the complement of \mathcal{U}^4 . It is an open submanifold of codimension 0.
- 3) *Non-stable axiumbilic Jets:* $(\mathcal{N}\mathcal{E})^4$, in the orbit of the axiumbilic jets for which:
 - $T = (\alpha_1\alpha_4 - \alpha_2\alpha_3)(r^2 + s^2) = 0$ or
 - $T \neq 0$ and conditions that characterize E_3 or E_4 axiumbilic points in Proposition 8 fail.

\mathcal{E}_{45}^1 is a closed variety of codimension 3, which can be expressed as the union of the following invariant strata:

- 3.1) *Non-Transversal jets:* \mathcal{E}_{45}^1 for which $T = 0$ and $\chi \neq 0$. It has codimension 3.
- 3.2) *Transversal-double jets:* $(\mathcal{E}_{34}^1)^4$, The Lie-Cartan field has a quadratic saddle-node in the projective line which is characterized by Proposition 11. It has codimension 3.
- 4) The *stable axiumbilic jets:* $\mathcal{U}\mathcal{E}^4$, the complement in \mathcal{U}^4 of $\mathcal{N}\mathcal{E}^4$.

Proposition 20. In the space of 1-parameter families of immersions, those whose 4-jet extension are transversal to the canonical axiumbilic stratification is open and dense.

Proof. Follows from Thom Transversality Theorem [6]. □

5. CONCLUDING COMMENTS

In this work was established the principal axial and the mean axial configurations in a neighborhood of the axiumbilic points of types E_{34}^1 and E_{45}^1 . The approach concerning methods and class of differentiability requirements is distinct from that presented in the work of Gutiérrez-Guñez-Castañeda in [3]. The use of the Lie-Cartan suspension method made possible the study of these points by means the classic theory of differential equations, in clear analogy with the saddle-node bifurcation of vector fields in the plane, following [1], [10] and [5].

The type E_{34}^1 satisfies the transversality condition of the curves a_0 and a_1 , Proposition 6, which amounts to the fact the Lie-Cartan surface remains regular in a neighborhood of the projective axis over the axiumbilic point. In this case there is a saddle-node equilibrium point of the Lie-Cartan vector field whose central separatrix is along the projective axis itself. The axial configurations are established in Proposition 10 and the qualitative change (bifurcation) between the types E_3 and E_4 , with the variation of a parameter in the space of immersions, is explained in Proposition 7. See Figure 10.

In the case E_{45}^1 the transversality condition fails, since curves a_0 and a_1 , Proposition 13, have quadratic contact at the axiumbilic point. Here the Lie-Cartan surface is not regular along the projective axis. It is established in Proposition 13 that there are four conic critical points of Morse type on the p -axis. At these points there are partially hyperbolic equilibria of the Lie-Cartan vector field. There is also a saddle-node equilibrium in the regular part of the surface whose central separatrix is transversal to the projective axis. The integral curves of the Lie - Cartan vector field on the regular components of the Lie - Cartan surface (which are four bi-punctured disks) are illustrated in Figure 12. Their projections on the plane give the axial configurations in a neighborhood of the axiumbilic point.

In Proposition 18 is established the one parameter variation (bifurcation) in the space of immersions. This leads to the fact that for small perturbations of an immersion with an axiumbilic point of this type it holds that two axiumbilic points, one of type E_4 and the other of type E_5 , bifurcate from E_{45}^1 or disappear leaving a neighborhood free from axiumbilic points, in full analogy with the saddle-node bifurcation [1] and [10]. See Figure 14.

In Proposition 20 the genericity of the points E_{34}^1 and E_{45}^1 is established in terms of stratification and transversality.

REFERENCES

- [1] Andronov, A. Gordon, I. Leontovich, E. Maier, G. Theory of Bifurcations of Dynamical Systems on a Plane. Israel Program for Scientific Translations, John Wiley, New York, 1973.
- [2] Gutiérrez, C., Guíñez, V. Simple Umbilic Points on Surfaces Immersed in \mathbb{R}^4 . Discrete Contin. Dyn. Syst. **9** (2003), 877-900. DOI: [10.3934/dcds.2003.9.877](https://doi.org/10.3934/dcds.2003.9.877)
- [3] Gutiérrez, C., Guíñez V., Castañeda, A. Quartic Differential Forms and Transversal Nets with Singularities. Discrete Contin. Dyn. Syst. **26** (2010), 225-249.
- [4] Gutiérrez, C., Guadalupe, I., Tribuzy, R., Guíñez, V. Lines of Curvature on Surfaces immersed in \mathbb{R}^4 . Bol. Soc. Bras. Mat., **28** (1997), 233-251. DOI: [10.1007/BF01233393](https://doi.org/10.1007/BF01233393)
- [5] Gutiérrez, C., Garcia, R., Sotomayor, J. Bifurcations of Umbilic Points and Related Principal Cycles. Journ. Dyn. and Diff. Eq., **16** (2004), 321-346.
- [6] H.T. Levine, Singularities of Differentiable Mappings, Lect. Notes in Math. **192**, 1971.
- [7] Little, J. A. On Singularities of Submanifolds of Higher Dimensional Euclidean Spaces. Ann. Mat. Pura Appl., **83** (1969), 261-335. DOI: [10.1007/BF02411172](https://doi.org/10.1007/BF02411172)
- [8] Mello, L. F. Mean Directionally Curved Lines on Surfaces Immersed in \mathbb{R}^4 . Publ. Mat., **47**,(2003), 415-440. DOI: [10.5565/PUBLMAT.47203.06](https://doi.org/10.5565/PUBLMAT.47203.06)
- [9] Mello, L. F. Orthogonal asymptotic lines on surfaces immersed in \mathbb{R}^4 . Rocky Mountain J. Math. **39** (2009), 1597-1612. DOI: [10.1216/RMJ-2009-39-5-1597](https://doi.org/10.1216/RMJ-2009-39-5-1597)
- [10] Sotomayor, J. Generic one-parameter families of vector fields on two-dimensional manifolds. Inst. Hautes Études Sci. Publ. Math. **43** (1974), 5-46. DOI: [10.1007/BF02684365](https://doi.org/10.1007/BF02684365)
- [11] Sotomayor, J., Garcia, R. Lines of Axial Curvature on Surfaces Immersed in \mathbb{R}^4 . Diff. Geom. and its Applications. **12** (2000), 253-269.
- [12] Sotomayor, J., Garcia, R. Differential Equations of Classical Geometry, a Qualitative Theory. Publicações Matemáticas. IMPA. 2009.

INSTITUTO DE MATEMÁTICA E ESTATÍSTICA, UNIVERSIDADE FEDERAL DE GOIÁS, CEP 74001-970, CAIXA POSTAL 131, GOIÂNIA, GOIÁS, BRAZIL

E-mail address: ragarcia@ufg.br

INSTITUTO DE MATEMÁTICA E ESTATÍSTICA, UNIVERSIDADE DE SÃO PAULO, RUA DO MATÃO 1010, CIDADE UNIVERITÁRIA, CEP 05508-090, SÃO PAULO, S. P, BRAZIL

E-mail address: sotp@ime.usp.br

INSTITUTO DE MATEMÁTICA E ESTATÍSTICA, UNIVERSIDADE DE SÃO PAULO, RUA DO MATÃO 1010, CIDADE UNIVERITÁRIA, CEP 05508-090, SÃO PAULO, S. P, BRAZIL

E-mail address: flausino@ime.usp.br